arXiv:math/0507511v3 [math.NT] 17 Aug 2005

A q-ANALOGUE OF LEHMER'S CONGRUENCE

Hao Pan

Abstract. We establish the q-analogue of a classical congruence of Lehmer. Also, the q-analogues of two congruences of Morley and Granville are given.

1. Introduction

In 1938, Lehmer [Leh] established an interesting congruence as follows:

$$\sum_{j=1}^{(p-1)/2}\frac{1}{j}\equiv -2\mathrm{Q}_p(2)+\mathrm{Q}_p(2)^2p \pmod{p^2}, \tag{1.1}$$

where $p\geq 3$ is prime and $\mathrm{Q}_p(2)=(2^{p-1}-1)/p$. Lehmer's congruence can be considered as an extension of the Wolstenholme's harmonic series congruence [W]

$$\sum_{j=1}^{p-1}\frac{1}{j}\equiv 0 \pmod{p^2}. \tag{1.2}$$

On the other hand, the q-analogues of some arithmetic congruences have been investigated by several authors (e.g., see [A], [F], [C], [GZ] and [PS]). Recently, Shi and Pan [SP] proved the following q-analogue of (1.2):

$$\sum_{j=1}^{p-1}\frac{1}{[j]_q}\equiv\frac{p-1}{2}(1-q)+\frac{p^2-1}{24}(1-q)^2[p]_q \pmod{[p]_q^2}, \tag{1.3}$$

where $[n]=(1-q^n)/(1-q)=1+q+\cdots+q^{n-1}$. Obviously (1.2) is deduced from (1.3) when $q\to 1$.

The main purpose of the present paper is to establish the q-analugue of Lehmer's congruence. Set

$$(a;q)_n=\begin{cases}(1-a)(1-aq)\cdots(1-aq^{n-1}) & \text{if } n\geq 1,\\ 1 & \text{if } n=0.\end{cases}$$

1991 *Mathematics Subject Classification.* Primary 11B65; Secondary 05A10, 05A30, 11A07.

It is easy to see that for any $m \geq 0$ with $p \nmid m$ we have a q-analogue of Fermat's little theorem

$$\frac{(q^m;q^m)_{p-1}}{(q;q)_{p-1}} \equiv 1 \pmod{[p]_q}. \tag{1.4}$$

Indeed, since

$$[m]_q = \frac{1-q^m}{1-q} \equiv \frac{1-q^n}{1-q} = [n]_q \pmod{[p]_q}$$

whenever $m \equiv n \pmod{p}$,

$$\frac{(q^m;q^m)_{p-1}}{(q;q)_{p-1}} = \prod_{j=1}^{p-1} \frac{1-q^{jm}}{1-q^j} = \prod_{j=1}^{p-1} \frac{[jm]_q}{[j]_q} \equiv 1 \pmod{[p]_q}.$$

So we can define the q-Fermat quotient by

$$\mathrm{Q}_p(m,q) = \frac{(q^m;q^m)_{p-1}/(q;q)_{p-1} - 1}{[p]_q}.$$

Theorem 1.1. *Let p be an odd prime. We have*

$$\begin{aligned} &2\sum_{j=1}^{(p-1)/2} \frac{1}{[2j]_q} + 2\mathrm{Q}_p(2,q) - \mathrm{Q}_p(2,q)^2[p]_q \\ \equiv&\left(\mathrm{Q}_p(2,q)(1-q) + \frac{p^2-1}{8}(1-q)^2\right)[p]_q \pmod{[p]_q^2}. \end{aligned} \tag{1.5}$$

In 1895, with the help of De Moivre's Theorem, Morley [M] proved that

$$(-1)^{\frac{p-1}{2}}\binom{p-1}{(p-1)/2} \equiv 4^{p-1} \pmod{p^3}. \tag{1.6}$$

for any prime $p \geq 5$. In [G1], Granville generalized the congruence of Morley and showed that

$$(-1)^{(p-1)(m-1)/2}\prod_{k=1}^{m-1}\binom{p-1}{\lfloor kp/m \rfloor} \equiv m^p - m + 1 \pmod{p^2}, \tag{1.7}$$

for any $m \geq 2$, where $\lfloor x \rfloor$ denotes the greatest integer not exceeding x. Now we can give the q-analogues of (1.6) and (1.7). For any $m, n \in \mathbb{N}$, define the q-binomial coefficients by

$$\begin{bmatrix} n \\ m \end{bmatrix}_q = \frac{(q;q)_n}{(q;q)_m(q;q)_{n-m}}$$

if $n \geq m$, and if $n < m$, then we let $\begin{bmatrix} n \\ m \end{bmatrix}_q = 0$. It is easy to see that $\begin{bmatrix} n \\ m \end{bmatrix}_q$ is a polynomial in q with integral coefficients, since the q-binomial coefficients satisfy the recurrence relation

$$\begin{bmatrix} n+1 \\ m \end{bmatrix}_q = q^m\begin{bmatrix} n \\ m \end{bmatrix}_q + \begin{bmatrix} n \\ m-1 \end{bmatrix}_q.$$

Theorem 1.2.

$$(-1)^{\frac{p-1}{2}}q^{\frac{p^2-1}{4}}\begin{bmatrix} p-1 \\ (p-1)/2 \end{bmatrix}_{q^2} \equiv (-q;q)_{p-1}^2 - \frac{p^2-1}{24}(1-q)^2[p]_q^2 \pmod{[p]_q^3} \quad (1.8)$$

for any prime $p \geq 5$.

Theorem 1.3. *Let $p \geq 5$ be a prime and $m \geq 2$ be an integer with $p \nmid m$. Then*

$$(-1)^{(p-1)(m-1)/2}q^M \prod_{k=1}^{m-1}\begin{bmatrix} p-1 \\ \lfloor kp/m \rfloor \end{bmatrix}_{q^m} \equiv \frac{m(q^m;q^m)_{p-1}}{(q;q)_{p-1}} - m + 1 \pmod{[p]_q^2}, \quad (1.9)$$

where

$$M = m\sum_{k=1}^{m-1}\binom{\lfloor kp/m \rfloor + 1}{2}.$$

The proofs of Theorems 1.1, 1.2 and 1.3 will be given in the next sections.

2. Some Lemmas

In this section we assume that p is a prime greater than 3. And the following lemmas will be used in the proofs of Theorems 1.1 and 1.2.

Lemma 2.1.

$$\sum_{j=1}^{p-1}\frac{1}{[j]_q} \equiv \frac{p-1}{2}(1-q) \pmod{[p]_q}, \quad (2.1)$$

$$\sum_{j=1}^{p-1}\frac{q^j}{[j]_q^2} \equiv -\frac{p^2-1}{12}(1-q)^2 \pmod{[p]_q} \quad (2.2)$$

and

$$\sum_{j=1}^{p-1}\frac{1}{[j]_q^2} \equiv -\frac{(p-1)(p-5)}{12}(1-q)^2 \pmod{[p]_q}. \quad (2.3)$$

Proof. See Theorem 4 in [A] and Lemma 2 in [SP].

Lemma 2.2.

$$q^{kp} = \sum_{j=0}^{k}(-1)^j\binom{k}{j}(1-q)^j[p]_q^j.$$

Proof.

$$\sum_{j=0}^{k}(-1)^j\binom{k}{j}(1-q)^j[p]_q^j = (1-(1-q)[p]_q)^k = (1-(1-q^p))^k = q^{kp}.$$

□

From Lemma 2.2, we deduce that

$$q^{kp} \equiv 1 - k(1-q)[p]_q + \frac{k(k-1)}{2}(1-q)^2[p]_q^2 \pmod{[p]_q^3}. \quad (2.4)$$

Lemma 2.3.

$$
\begin{aligned}
&4\sum_{1\le j<k\le p-1}\frac{(-1)^k}{[j]_q[k]_q}\\
\equiv&\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\bigg)^2+(p-3)(1-q)\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}+\frac{(p-1)(p+7)}{12}(1-q)^2 \pmod{[p]_q}.
\end{aligned}
\tag{2.5}
$$

Proof. Since p is odd,

$$
\begin{aligned}
\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\bigg)^2=&\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\bigg)\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}-(1-q)\sum_{j=1}^{p-1}(-1)^j\bigg)\\
=&\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\bigg)\bigg(\sum_{j=1}^{p-1}\frac{(-q)^j}{[j]_q}\bigg)\\
=&\sum_{k=2}^{2p-2}(-1)^k\sum_{j=\max\{1,k-p+1\}}^{\min\{k-1,p-1\}}\frac{q^j}{[j]_q[k-j]_q}.
\end{aligned}
$$

Then we have

$$
\begin{aligned}
&\bigg(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\bigg)\bigg(\sum_{j=1}^{p-1}\frac{(-q)^j}{[j]_q}\bigg)-(-1)^p\sum_{j=1}^{p-1}\frac{q^j}{[j]_q[p-j]_q}\\
=&\sum_{k=2}^{p-1}(-1)^k\sum_{j=1}^{k-1}\frac{q^j}{[j]_q[k-j]_q}+\sum_{k=p+1}^{2p-2}(-1)^k\sum_{j=k-p+1}^{p-1}\frac{q^j}{[j]_q[k-j]_q}\\
=&\sum_{k=2}^{p-1}(-1)^k\sum_{j=1}^{k-1}\frac{q^j}{[j]_q[k-j]_q}+\sum_{l=2}^{p-1}(-1)^{2p-l}\sum_{j=p-l+1}^{p-1}\frac{q^j}{[j]_q[2p-l-j]_q}\\
&\text{(here } l=2p-k)\\
=&\sum_{k=2}^{p-1}(-1)^k\sum_{j=1}^{k-1}\frac{q^j}{[j]_q[k-j]_q}+\sum_{l=2}^{p-1}(-1)^l\sum_{i=1}^{l-1}\frac{q^{p+i-l}}{[p+i-l]_q[p-i]_q}.\\
&\text{(here } i=l+j-p)
\end{aligned}
$$

Note that

$$
\frac{q^{p+i-l}}{[p+i-l]_q[p-i]_q}=\frac{q^{p+i-l}(1-q)^2}{(1-q^{p+i-l})(1-q^{p-i})}\equiv\frac{q^i(1-q)^2}{(1-q^{l-i})(1-q^i)} \pmod{[p]_q}.
$$

It follows that

$$\begin{aligned}
&\left(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\right)\left(\sum_{j=1}^{p-1}\frac{(-q)^j}{[j]_q}\right)+\sum_{j=1}^{p-1}\frac{q^j}{[j]_q[p-j]_q}\\
\equiv&\sum_{k=2}^{p-1}(-1)^k\sum_{j=1}^{k-1}\frac{q^j}{[j]_q[k-j]_q}+\sum_{l=2}^{p-1}(-1)^l\sum_{i=1}^{l-1}\frac{q^i}{[i]_q[l-i]_q}\\
=&2\sum_{k=2}^{p-1}(-1)^k\sum_{j=1}^{k-1}\frac{q^j(1-q)^2}{(1-q^j)(1-q^{k-j})}\\
=&2\sum_{k=2}^{p-1}\frac{(-1)^kq^k(1-q)^2}{1-q^k}\sum_{j=1}^{k-1}\left(\frac{1}{q^{k-j}-q^k}+\frac{1}{1-q^{k-j}}\right)\\
=&2\sum_{1\le j<k\le p-1}\frac{(-1)^k(q^k+q^j)}{[j]_q[k]_q}\pmod{[p]_q}.
\end{aligned}$$

We can write

$$\begin{aligned}
&\sum_{1\le j<k\le p-1}\frac{(-1)^k(q^k+q^j)}{[j]_q[k]_q}\\
=&\sum_{1\le j<k\le p-1}\frac{(-1)^k(2-(1-q^k)-(1-q^j))}{[j]_q[k]_q}\\
=&2\sum_{1\le j<k\le p-1}\frac{(-1)^k}{[j]_q[k]_q}-(1-q)\left(\sum_{k=2}^{p-1}\frac{(-1)^k(k-1)}{[k]_q}+\sum_{j=1}^{p-2}\frac{1}{[j]_q}\sum_{k=j+1}^{p-1}(-1)^k\right).
\end{aligned}$$

Now

$$\begin{aligned}
\sum_{k=2}^{p-1}\frac{(-1)^k(k-1)}{[k]_q}=&\frac{1}{2}\sum_{k=1}^{p-1}\left(\frac{(-1)^k(k-1)}{[k]_q}+\frac{(-1)^{p-k}(p-k-1)}{[p-k]_q}\right)\\
=&\frac{1}{2}\sum_{k=1}^{p-1}(-1)^k(k-1)\left(\frac{1}{[k]_q}+\frac{1}{[p-k]_q}\right)+\frac{p-2}{2}\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\\
=&\frac{1}{2}\sum_{k=1}^{p-1}(-1)^k(k-1)\left(\frac{[p]_q}{[k][p-k]_q}+(1-q)\right)+\frac{p-2}{2}\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\\
\equiv&\frac{p-1}{4}(1-q)+\frac{p-2}{2}\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\pmod{[p]_q}.
\end{aligned}$$

And from (2.1) we have

$$\begin{aligned}\sum_{j=1}^{p-2}\frac{1}{[j]_q}\sum_{k=j+1}^{p-1}(-1)^k =&\frac{1}{2}\sum_{j=1}^{p-1}\frac{1-(-1)^j}{[j]_q}\\ \equiv&\frac{(p-1)}{4}(1-q)-\frac{1}{2}\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q} \pmod{[p]_q}.\end{aligned}$$

Finally by (2.3),

$$\begin{aligned}\sum_{j=1}^{p-1}\frac{q^j}{[j]_q[p-j]_q}=\sum_{j=1}^{p-1}\frac{q^{p-j}}{[j]_q[p-j]_q}=&\sum_{j=1}^{p-1}\frac{q^p}{[j]_q([p]-[j]_q)}\\ \equiv&\frac{(p-1)(p-5)}{12}(1-q)^2 \pmod{[p]_q}.\end{aligned}$$

Thus combining the equations and congruences above, we obtain that

$$\begin{aligned}&4\sum_{1\le j<k\le p-1}(-1)^k\frac{1}{[j]_q[k]_q}-\left(\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\right)\left(\sum_{j=1}^{p-1}\frac{(-q)^j}{[j]_q}\right)\\ \equiv&(p-3)(1-q)\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}+\frac{(p-1)(p+7)}{12}(1-q)^2 \pmod{[p]_q}.\end{aligned}$$

□

Lemma 2.4.

$$\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}\equiv 2\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}-\frac{p-1}{2}(1-q)-\frac{p^2-1}{24}(1-q)^2[p]_q \pmod{[p]_q^2}. \quad (2.6)$$

Proof. Clearly

$$\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}=\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}-\sum_{j=1}^{(p-1)/2}\frac{1}{[2j-1]_q}=\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}-\sum_{j=1}^{(p-1)/2}\frac{1}{[p-2j]_q}.$$

Observe that

$$\frac{1}{[p-2j]_q}=\frac{q^{2j}}{[p]_q-[2j]_q}=\frac{q^{2j}([p]_q+[2j]_q)}{[p]_q^2-[2j]_q^2}\equiv-\frac{q^{2j}([p]_q+[2j]_q)}{[2j]_q^2} \pmod{[p]_q^2}.$$

And by (2.2), we have

$$\begin{aligned}-\frac{p^2-1}{12}(1-q)^2\equiv\sum_{j=1}^{p-1}\frac{q^j}{[j]_q^2}&=\sum_{j=1}^{(p-1)/2}\frac{q^{2j}}{[2j]_q^2}+\sum_{j=1}^{(p-1)/2}\frac{q^{p-2j}}{[p-2j]_q^2}\\&=\sum_{j=1}^{(p-1)/2}\frac{q^{2j}}{[2j]_q^2}+\sum_{j=1}^{(p-1)/2}\frac{q^{p+2j}}{([p]_q-[2j]_q)^2}\\&\equiv 2\sum_{j=1}^{(p-1)/2}\frac{q^{2j}}{[2j]_q^2}\ (\mathrm{mod}\ [p]_q).\end{aligned}\tag{2.7}$$

Hence

$$\begin{aligned}\sum_{j=1}^{p-1}\frac{(-1)^j}{[j]_q}&\equiv\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}+\sum_{j=1}^{(p-1)/2}\frac{q^{2j}([p]_q+[2j]_q)}{[2j]_q^2}\\&\equiv\sum_{j=1}^{(p-1)/2}\frac{1+q^{2j}}{[2j]_q}-\frac{p^2-1}{24}(1-q)^2[p]_q\\&=2\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}-\frac{p-1}{2}(1-q)-\frac{p^2-1}{24}(1-q)^2[p]_q\ (\mathrm{mod}\ [p]_q^2).\end{aligned}$$

□

3. Proofs of Theorems 1.1 and 1.2

Proof of Theorem 1.1. One can directly verify (1.5) when $p=3$. So below we assume that $p\geq 5$. It is well-known (cf. Corollary 10.2.2 of [AAR]) that

$$(x;q)_n=\sum_{j=0}^{n}\begin{bmatrix}n\\j\end{bmatrix}_q q^{\binom{j}{2}}(-x)^j.$$

Then we have

$$\begin{aligned}\frac{(-1;q)_q-q^{\binom{p}{2}}-1}{[p]_q}&=\frac{1}{[p]_q}\sum_{k=1}^{p-1}\begin{bmatrix}p\\k\end{bmatrix}_q q^{\binom{k}{2}}\\&=\sum_{k=1}^{p-1}\frac{1}{[k]_q}\prod_{j=1}^{k-1}\frac{q^j(1-q^{p-j})}{1-q^j}\\&=\sum_{k=1}^{p-1}\frac{1}{[k]_q}\prod_{j=1}^{k-1}\left(\frac{[p]_q}{[j]_q}-1\right)\\&\equiv[p]_q\sum_{1\leq j<k\leq p-1}\frac{(-1)^k}{[j]_q[k]_q}-\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\ (\mathrm{mod}\ [p]_q^2).\end{aligned}\tag{3.1}$$

Consequently

$$\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\equiv-\frac{(-1;q)_q-q^{\binom{p}{2}}-1}{[p]_q}\equiv-\frac{2(-q;q)_{q-1}-2}{[p]_q}-\frac{p-1}{2}(1-q)\ (\mathrm{mod}\ [p]_q).$$

Thus applying Lemma 2.3, we have

$$\begin{aligned}&\sum_{1\le j<k\le p-1}\frac{(-1)^k}{[j]_q[k]_q}-\frac{(p-1)(p+7)}{48}(1-q)^2\\ \equiv&\frac{1}{4}\bigg(\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}\bigg)\bigg(\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}+(p-3)(1-q)\bigg)\\ \equiv&\frac{1}{4}\bigg(-2\mathrm{Q}_p(2,q)-\frac{p-1}{2}(1-q)\bigg)\bigg(-2\mathrm{Q}_p(2,q)+\frac{p-5}{2}(1-q)\bigg)\\ =&\mathrm{Q}_p(2,q)^2+\mathrm{Q}_p(2,q)(1-q)-\frac{(p-1)(p-5)}{16}(1-q)^2\ (\mathrm{mod}\ [p]_q).\end{aligned}\tag{3.2}$$

On the other hand, it follows from (2.4) that

$$\begin{aligned}&\frac{(-1;q)_q-q^{\binom{p}{2}}-1}{[p]_q}\\ \equiv&\frac{2(-q;q)_{q-1}-2}{[p]_q}+\frac{p-1}{2}(1-q)-\frac{(p-1)(p-3)}{8}(1-q)^2[p]_q\ (\mathrm{mod}\ [p]_q^2).\end{aligned}$$

Then by Lemma 2.4,

$$\begin{aligned}&\sum_{k=1}^{p-1}\frac{(-1)^k}{[k]_q}+\frac{(-1;q)_q-q^{\binom{p}{2}}-1}{[p]_q}\\ \equiv&2\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}+2\mathrm{Q}_p(2,q)-\frac{(p-1)(p-2)}{6}(1-q)^2[p]_q\ (\mathrm{mod}\ [p]_q^2).\end{aligned}\tag{3.3}$$

Combining (3.1), (3.2) and (3.3), the desired (1.5) is obtained. □

Proof of Theorem 1.2. Since

$$\begin{bmatrix}p-1\\(p-1)/2\end{bmatrix}_{q^2}=\prod_{j=1}^{(p-1)/2}\frac{[p-j]_{q^2}}{[j]_{q^2}}=\prod_{j=1}^{(p-1)/2}\frac{[p]_{q^2}-[j]_{q^2}}{q^{2j}[j]_{q^2}},$$

we have

$$\begin{aligned}&(-1)^{\frac{p-1}{2}}q^{\frac{p^2-1}{4}}\begin{bmatrix}p-1\\(p-1)/2\end{bmatrix}_{q^2}\\ =&\prod_{j=1}^{(p-1)/2}\bigg(1-\frac{[p]_{q^2}}{[j]_{q^2}}\bigg)\\ \equiv&1-\frac{1+q^p}{1+q}\sum_{j=1}^{(p-1)/2}\frac{[p]_q}{[j]_{q^2}}+\frac{(1+q^p)^2}{(1+q)^2}\sum_{1\le j<k\le(p-1)/2}\frac{[p]_q^2}{[j]_{q^2}[k]_{q^2}}\ (\mathrm{mod}\ [p]_{q^2}^3).\end{aligned}\tag{3.4}$$

From Theorem 1.1, we deduce that

$$\begin{aligned}&\frac{1+q^p}{1+q}\sum_{j=1}^{(p-1)/2}\frac{1}{[j]_{q^2}}\\ =&(1+q^p)\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}\\ \equiv&-(1+q^p)\mathrm{Q}_p(2,q)+\frac{1+q^p}{2}\left(\mathrm{Q}_p(2,q)^2+\mathrm{Q}_p(2,q)(1-q)+\frac{p^2-1}{8}(1-q)^2\right)[p]_q\\ \equiv&-(1+q^p)\mathrm{Q}_p(2,q)+\mathrm{Q}_p(2,q)^2[p]_q+\mathrm{Q}_p(2,q)(1-q)[p]_q+\frac{p^2-1}{8}(1-q)^2[p]_q\\ &\pmod{[p]_q^2}.\end{aligned}\tag{3.5}$$

Notice that

$$\begin{aligned}\sum_{1\le j<k\le(p-1)/2}\frac{1}{[j]_{q^2}[k]_{q^2}}=&\frac{1}{2}\left(\left(\sum_{j=1}^{(p-1)/2}\frac{1}{[j]_{q^2}}\right)^2-\sum_{j=1}^{(p-1)/2}\frac{1}{[j]_{q^2}^2}\right)\\ =&\frac{(1+q)^2}{2}\left(\left(\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}\right)^2-\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q^2}\right).\end{aligned}$$

And Theorem 1.1 implies that

$$\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}\equiv-\mathrm{Q}_p(2,q)\pmod{[p]_q}.$$

Then using (2.7),

$$\begin{aligned}\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q^2}=&\sum_{j=1}^{(p-1)/2}\frac{q^{2j}}{[2j]_q^2}+(1-q)\sum_{j=1}^{(p-1)/2}\frac{1}{[2j]_q}\\ \equiv&-\frac{p^2-1}{24}(1-q)^2-\mathrm{Q}_p(2,q)(1-q)\pmod{[p]_q}.\end{aligned}$$

Consequently

$$\begin{aligned}&\frac{2}{(1+q)^2}\sum_{1\le j<k\le(p-1)/2}\frac{1}{[j]_{q^2}[k]_{q^2}}\\ \equiv&\mathrm{Q}_p(2,q)^2+\mathrm{Q}_p(2,q)(1-q)+\frac{p^2-1}{24}(1-q)^2\pmod{[p]_q}.\end{aligned}\tag{3.6}$$

Thus it follows from (3.4), (3.5) and (3.6) that

$$\begin{aligned}&(-1)^{\frac{p-1}{2}}q^{\frac{p^2-1}{4}}\begin{bmatrix}p-1\\(p-1)/2\end{bmatrix}_{q^2}-1\\ \equiv&[p]_q^2\cdot\frac{4}{(1+q)^2}\sum_{1\le j<k\le(p-1)/2}\frac{1}{[j]_{q^2}[k]_{q^2}}-[p]_q\cdot\frac{1+q^p}{1+q}\sum_{j=1}^{(p-1)/2}\frac{1}{[j]_{q^2}}\\ \equiv&\mathrm{Q}_p(2,q)^2[p]_q^2+(1+q^p)\mathrm{Q}_p(2,q)[p]_q+\mathrm{Q}_p(2,q)(1-q)[p]_q^2-\frac{p^2-1}{24}(1-q)^2[p]_q^2\\ =&((-q;q)_{p-1}-1)((-q;q)_{p-1}+1)-\frac{p^2-1}{24}(1-q)^2[p]_q^2 \pmod{[p]_q^3}.\end{aligned}$$

□

4. Fermat Quotient

Lemma 4.1. *Let p be an odd prime. Suppose that m is a positive integer with $(m,p)=1$. Then*

$$\mathrm{Q}_p(m,q)\equiv\sum_{j=1}^{p-1}\frac{\lfloor jm/p\rfloor}{[jm]_q}-\frac{(p-1)(m-1)}{2}(1-q) \pmod{[p]_q}. \tag{4.1}$$

Proof. For each $j\in\{1,2,\ldots,p-1\}$, let

$$r_j=jm-\lfloor jm/p\rfloor p.$$

Then

$$\begin{aligned}\frac{(q^m;q^m)_{p-1}}{(q;q)_{p-1}}=\prod_{j=1}^{p-1}\frac{1-q^{jm}}{1-q^j}=&\prod_{j=1}^{p-1}\left(\frac{1-q^{r_j}}{1-q^j}+\frac{q^{r_j}(1-q^{\lfloor jm/p\rfloor p})}{1-q^j}\right)\\ =&\prod_{j=1}^{p-1}\frac{1-q^{r_j}}{1-q^j}\left(1+\frac{q^{r_j}(1-q^{\lfloor jm/p\rfloor p})}{1-q^{r_j}}\right)\end{aligned}$$

Since r_j runs through $1,2,\ldots,p-1$ as j does so, we have

$$\begin{aligned}\frac{(q^m;q^m)_{p-1}}{(q;q)_{p-1}}=&\prod_{j=1}^{p-1}\left(1+\frac{q^{r_j}(1-q^{\lfloor jm/p\rfloor p})}{1-q^{r_j}}\right)\\ \equiv&1+(1-q^p)\sum_{j=1}^{p-1}\frac{q^{r_j}}{1-q^{r_j}}\cdot\frac{1-q^{\lfloor jm/p\rfloor p}}{1-q^p}\\ \equiv&1+(1-q^p)\sum_{j=1}^{p-1}\left\lfloor\frac{jm}{p}\right\rfloor\frac{q^{r_j}}{1-q^{r_j}}\\ \equiv&1+[p]_q\sum_{j=1}^{p-1}\left\lfloor\frac{jm}{p}\right\rfloor\frac{q^{jm}}{[jm]_q} \pmod{[p]_q^2}.\end{aligned}$$

Finally,

$$\begin{aligned}\sum_{j=1}^{p-1}\left\lfloor\frac{jm}{p}\right\rfloor\frac{q^{jm}}{[jm]_q}=&\sum_{j=1}^{p-1}\frac{\lfloor jm/p\rfloor}{[jm]_q}-(1-q)\sum_{j=1}^{p-1}\left\lfloor\frac{jm}{p}\right\rfloor\\=&\sum_{j=1}^{p-1}\frac{\lfloor jm/p\rfloor}{[jm]_q}-\frac{(p-1)(m-1)}{2}(1-q).\end{aligned}$$

We are done. □

Remark. Letting $q\to 1$ in (4.1), we obtain that

$$\frac{m^p-m}{p}\equiv\sum_{j=1}^{p-1}\frac{\lfloor jm/p\rfloor}{j}\pmod p,$$

which was firstly discovered by Lerch [Ler].

Proof of Theorem 1.3. We write

$$\left[\begin{matrix}p-1\\ \lfloor kp/m\rfloor\end{matrix}\right]_{q^m}=\prod_{j=1}^{\lfloor kp/m\rfloor}\frac{[p]_{q^m}-[j]_{q^m}}{q^{jm}[j]_{q^m}}=q^{-m\binom{\lfloor kp/m\rfloor+1}{2}}\prod_{j=1}^{\lfloor kp/m\rfloor}\left(\frac{[p]_{q^m}}{[j]_{q^m}}-1\right).$$

As $p\nmid m$, $[p]_q$ divides $[p]_{q^m}=(1-q^{mp})/(1-q^m)$. Thus

$$\begin{aligned}&(-1)^{(p-1)(m-1)/2}q^{\sum_{k=1}^{m-1}m\binom{\lfloor kp/m\rfloor+1}{2}}\prod_{k=1}^{m-1}\left[\begin{matrix}p-1\\ \lfloor kp/m\rfloor\end{matrix}\right]_{q^m}\\=&\prod_{k=1}^{m-1}\prod_{j=1}^{\lfloor kp/m\rfloor}\left(1-\frac{[p]_{q^m}}{[j]_{q^m}}\right)\\\equiv&1-[p]_{q^m}\sum_{k=1}^{m-1}\sum_{1\le j<kp/m}\frac{1}{[j]_{q^m}}\\&\text{(here } kp/m\notin\mathbb{Z},\ \text{so } j\le\lfloor kp/m\rfloor<kp/m)\\=&1-[p]_{q^m}\sum_{j=1}^{p-1}\frac{m-1-\lfloor jm/p\rfloor}{[j]_{q^m}}\pmod{[p]_q^2}.\end{aligned}$$

In view of (2.1) and Lemma 4.1, we have

$$\begin{aligned}&[p]_{q^m}\sum_{j=1}^{p-1}\frac{m-1-\lfloor jm/p\rfloor}{[j]_{q^m}}\\=&(m-1)[mp]_q\sum_{j=1}^{p-1}\frac{1}{[jm]_q}-[mp]_q\sum_{j=1}^{p-1}\frac{\lfloor jm/p\rfloor}{[jm]_q}\\\equiv&(m-1)[mp]_q\cdot\frac{(p-1)}{2}(1-q)-[mp]_q\mathrm{Q}_p(m,q)-\frac{(p-1)(m-1)}{2}(1-q)[mp]_q\\\equiv&-m[p]_q\mathrm{Q}_p(m,q)\pmod{[p]_q^2}.\end{aligned}$$

This concludes our proof. $\square$

Remark. For further developments of Granville's congruence (1.7), the reader is referred to [S].

5. A conjecture of Skula

Recently with help of polynomials over finite fields, Granville [G2] confirmed a conjecture of Skula:

$$\left(\frac{2^{p-1}-1}{p}\right)^2 \equiv -\sum_{j=1}^{p-1}\frac{2^j}{j^2} \pmod{[p]_q} \tag{5.1}$$

for any prime $p \geq 5$. Using our q-analogue of Lehmer's congruence, we have the following q-analogue of (5.1):

Theorem 5.1. *Let $p \geq 5$ be a prime. Then*

$$\begin{aligned}&\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q^2}+\mathrm{Q}_p(2,q)^2\\ \equiv &-(p-1)\mathrm{Q}_p(2,q)(1-q)-\frac{(7p-5)(p-1)}{24}(1-q)^2 \pmod{[p]_q}.\end{aligned} \tag{5.2}$$

Lemma 5.2.

$$\sum_{k=0}^{n}(-1)^k\begin{bmatrix}n\\k\end{bmatrix}_q q^{\binom{n-k}{2}}(-q;q)_k=(-1)^n q^{\binom{n+1}{2}}. \tag{5.3}$$

Proof. From the well-known q-binomial theorem (cf. Theorem 10.2.1 of [AAR]), we have

$$\sum_{k=0}^{\infty}\frac{(-1)^k q^{\binom{k}{2}}}{(q;q)_k}x^k=(x;q)_\infty$$

and

$$\sum_{k=0}^{\infty}\frac{(-q;q)_k}{(q;q)_k}x^k=\frac{(-qx;q)_\infty}{(x;q)_\infty}.$$

Then by comparing the coefficient of x^n in the both sides of

$$(x;q)_\infty\cdot\frac{(-qx;q)_\infty}{(x;q)_\infty}=(-qx;q)_\infty,$$

we obtain that

$$\sum_{k=0}^{n}\frac{(-1)^{n-k}q^{\binom{n-k}{2}}(-q;q)_k}{(q;q)_{n-k}(q;q)_k}=\frac{q^{\binom{n}{2}+n}}{(q;q)_n},$$

which is an equivalent form of (5.3). $\square$

Corollary 5.3. *For any odd prime p, we have*

$$\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q}\equiv -2\mathrm{Q}_p(2,q)-(p-1)(1-q)\ (\mathrm{mod}\ [p]_q). \tag{5.4}$$

Proof. From Lemma 5.2, we deduce that

$$\begin{aligned}\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q}\equiv&\sum_{j=1}^{p-1}\frac{q^{p(p-1)/2-jp+j}(-q;q)_j}{[j]_q}\\ \equiv&-\frac{1}{[p]_q}\sum_{j=1}^{p-1}(-1)^j q^{\binom{p}{2}+\binom{j}{2}-jp+j}\begin{bmatrix}p\\ j\end{bmatrix}_q(-q;q)_j\\ =&-\frac{1}{[p]_q}\sum_{j=1}^{p-1}(-1)^j q^{\binom{p-j}{2}}\begin{bmatrix}p\\ j\end{bmatrix}_q(-q;q)_j\\ =&-\frac{1}{[p]_q}((-1)^p q^{\binom{p+1}{2}}-q^{\binom{p}{2}}-(-1)^p(-q;q)_p)\ (\mathrm{mod}\ [p]_q).\end{aligned}$$

Notice that

$$\frac{2-q^{\binom{p+1}{2}}-q^{\binom{p}{2}}}{[p]_q}\equiv\frac{p+1}{2}(1-q)+\frac{p-1}{2}(1-q)=p(1-q)\ (\mathrm{mod}\ [p]_q),$$

and that

$$\begin{aligned}\frac{(-q;q)_p-2}{[p]_q}=\frac{(-q;q)_{p-1}(1+q^p)-2}{[p]_q}=&\frac{2(-q;q)_{p-1}-2}{[p]_q}-(1-q)(-q;q)_{p-1}\\ \equiv&2\mathrm{Q}_p(2,q)-(1-q)\ (\mathrm{mod}\ [p]_q).\end{aligned}$$

Hence

$$\begin{aligned}\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q}\equiv&-\frac{(-q;q)_p-q^{\binom{p+1}{2}}-q^{\binom{p}{2}}}{[p]_q}\\ \equiv&-2\mathrm{Q}_p(2,q)-(p-1)(1-q)\ (\mathrm{mod}\ [p]_q).\end{aligned}$$

We are done. □

Remark. Corollary 5.3 is the q-analogue of an observation of Glashier:

$$\frac{2^{p-1}-1}{p}\equiv-\sum_{j=1}^{p-1}\frac{2^{j-1}}{j}\ (\mathrm{mod}\ p). \tag{5.5}$$

Lemma 5.4.

$$\sum_{k=1}^{n}(-1)^k \begin{bmatrix} n\\ k\end{bmatrix}_q q^{\binom{n-k}{2}}\frac{(-q;q)_k}{[k]_q} = q^{\binom{n}{2}}\sum_{k=1}^{n}\frac{(-q)^k-1}{[k]_q}. \tag{5.6}$$

Proof. We make an induction on n. The case $n = 1$ is trivial. Assume that $n > 1$ and that (5.6) holds for the smaller values of n. Then we conclude that

$$\begin{aligned}
&\sum_{k=1}^{n}(-1)^k \begin{bmatrix} n\\ k\end{bmatrix}_q q^{\binom{n-k}{2}}\frac{(-q;q)_k}{[k]_q}\\
=&\sum_{k=1}^{n}(-1)^k\left(q^k\begin{bmatrix} n-1\\ k\end{bmatrix}_q + \begin{bmatrix} n-1\\ k-1\end{bmatrix}_q\right)q^{\binom{n-k}{2}}\frac{(-q;q)_k}{[k]_q}\\
=&q^{n-1}\sum_{k=1}^{n-1}(-1)^k\begin{bmatrix} n-1\\ k\end{bmatrix}_q q^{\binom{n-k-1}{2}}\frac{(-q;q)_k}{[k]_q} + \frac{1}{[n]_q}\sum_{k=1}^{n}(-1)^k\begin{bmatrix} n\\ k\end{bmatrix}_q q^{\binom{n-k}{2}}(-q;q)_k\\
=&q^{\binom{n}{2}}\sum_{k=1}^{n-1}\frac{(-q)^k-1}{[k]_q} + \frac{1}{[n]_q}((-1)^n q^{\binom{n+1}{2}} - q^{\binom{n}{2}}),
\end{aligned}$$

where in the last step we apply the induction hypothesis and Lemma 5.2. □

Proof of Theorem 5.1. Using Lemma 5.4, we have

$$\begin{aligned}
\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q^2} \equiv&\sum_{j=1}^{p-1}\frac{q^{p(p-1)/2-jp+j}(-q;q)_j}{[j]_q^2}\\
\equiv&-\frac{1}{[p]_q}\sum_{j=1}^{p-1}(-1)^j q^{\binom{p-j}{2}}\begin{bmatrix} p\\ j\end{bmatrix}_q\frac{(-q;q)_j}{[j]_q}\\
=&-\frac{q^{\binom{p}{2}}}{[p]_q}\sum_{j=1}^{p}\frac{(-q)^j-1}{[j]_q} + (-1)^p\frac{(-q;q)_p}{[p]_q^2}\\
=&-\frac{q^{\binom{p}{2}}}{[p]_q}\sum_{j=1}^{p-1}\frac{(-q)^j-1}{[j]_q} - \frac{(-q;q)_p - q^{\binom{p+1}{2}} - q^{\binom{p}{2}}}{[p]_q^2} \pmod{[p]_q}.
\end{aligned}$$

With help of (1.3) and Theorem 1.1,

$$\begin{aligned}\sum_{k=1}^{p-1}\frac{(-q)^j-1}{[j]_q} =& -\sum_{k=1}^{p-1}\frac{(-1)^j(1-q^j)}{[j]_q}+\sum_{k=1}^{p-1}\frac{(-1)^j-1}{[j]_q}\\ =& -\sum_{j=1}^{(p-1)/2}\frac{2}{[2j-1]_q}\\ =& \sum_{j=1}^{(p-1)/2}\frac{2}{[2j]_q}-\sum_{k=1}^{p-1}\frac{2}{[j]_q}\\ \equiv& -2\mathrm{Q}_p(2,q)+\mathrm{Q}_p(2,q)^2[p]_q+\mathrm{Q}_p(2,q)(1-q)[p]_q\\ &+\frac{p^2-1}{8}(1-q)^2[p]_q-((p-1)(1-q)+\frac{p^2-1}{12}(1-q)^2[p]_q)\\ &\pmod{[p]_q^2}.\end{aligned}$$

And by (2.4) we have

$$\begin{aligned}&\frac{(-q;q)_p-q^{\binom{p+1}{2}}-q^{\binom{p}{2}}}{[p]_q^2}\\ =&\frac{2(-q;q)_{p-1}-q^{\binom{p+1}{2}}-q^{\binom{p}{2}}}{[p]_q^2}-\frac{(-q;q)_{p-1}}{[p]_q}(1-q)\\ \equiv&\frac{2(-q;q)_{p-1}-2}{[p]_q^2}+\frac{p}{[p]_q}(1-q)-\frac{(p-1)^2}{4}(1-q)^2-\frac{(-q;q)_{p-1}}{[p]_q}(1-q)\\ =&\frac{2\mathrm{Q}_p(2,q)}{[p]_q}+\frac{(p-1)(1-q)}{[p]_q}-\frac{(p-1)^2}{4}(1-q)^2-\mathrm{Q}_p(2,q)(1-q)\pmod{[p]_q}.\end{aligned}$$

Therefore

$$\begin{aligned}&\sum_{j=1}^{p-1}\frac{q^j(-q;q)_j}{[j]_q^2}\\ \equiv&\left(\frac{(p-1)(1-q)}{2}-\frac{1}{[p]_q}\right)\sum_{j=1}^{p-1}\frac{(-q)^j-1}{[j]_q}-\frac{(-q;q)_p-q^{\binom{p+1}{2}}-q^{\binom{p}{2}}}{[p]_q^2}\\ \equiv&-(p-1)\mathrm{Q}_p(2,q)(1-q)-\mathrm{Q}_p(2,q)^2-\frac{(7p-5)(p-1)}{24}(1-q)^2\pmod{[p]_q}.\end{aligned}$$

□

Acknowledgment. I thank my advisor, Prof. Zhi-Wei Sun, for his useful comments on this paper.

References

- [A] G. E. Andrews, *q-Analogs of the binomial coefficient congruences of Babbage, Wolstenholme and Glaisher*, Discrete Math. **204** (1999), 15-25.
- [AAR] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.
- [C] W. E. Clark, *q-analog of a binomial coefficient congruence*, Internat. J. Math. and Math. Sci. **18**, 197-200.
- [F] R. D. Fray, *Congruence properties of ordinary and q-binomial coefficients*, Duke Math. J. **34** (1967), 467-480.
- [G1] A. Granville, *Arithmetic Properties of Binomial Coefficients I: Binomial coefficients modulo prime powers,* in Organicmathematics (Burnady,BC,1995), CMS Conf. Proc., 20, Amer. Math. Soc., Providence, RI, 1997, pp. 253-275.
- [G2] A. Granville, *The Square of the Fermat Quotient*, Integers **3** (2003), A8.
- [GZ] Victor J. W. Guo and J. Zeng, *Some arithmetic properties of the q-Euler numbers and q-Salié numbers*, European J. Combin., to appear, arXiv:math.CO/0504569.
- [Leh] E. Lehmer, *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Annals of Math. **39** (1938), 350-360.
- [Ler] M. Lerch, *Zur Theorie des Fermatschen Quotienten $(a^{p-1}-1)/p = q(a)$*, Math. Ann. **60** (1905), 471-490.
- [M] F. Morley, *Note on the congruence $2^{4n} \equiv (-1)^n (2n)!/(n!)^2$, where $2n+1$ is a prime*, Annals of Math. **9** (1895), 168-170.
- [PS] H. Pan and Z.-W. Sun, *On q-Euler numbers, q-Salié numbers and q-Carlitz numbers*, preprint, arXiv:math.CO/0505548.
- [S] Z.-W. Sun, *Products of binomial coefficients modulo p^2*, Acta Arith. **97** (2001), 87-98.
- [SP] L.-L. Shi and H. Pan, *A q-analogue of Wolstenholme's harmonic series congruence*, preprint, arXiv:math.NT/0507495.
- [W] J. Wolstenholme, *On certain properties of prime numbers*, Quart J. Math. **5** (1862), 35-39.

Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

E-mail address: `haopan79@yahoo.com.cn`